

References

Jayaram P, Kennedy DJ, Yeh P, and Dragoo J. Chondrotoxic Effects of Local Anesthetics on Human Knee Articular Cartilage: A Systematic Review. *PMR*. 2019;11(4):379-400.

Raymond HE, Barbera JP, Shah KC, Zubizarreta N, Huang HH, Poeran J, Chen DD, Moucha CS, and Hayden BL. Risk of Infection After Total Knee or Hip Arthroplasty After Receipt of Multiple Corticosteroid or Hyaluronic Acid Injections. *J Am Acad Orthop Surg*. 2023;31(19): e868-e875.

Baums MH, Aquilina J, Pérez-Prieto D, Sleiman O, Geropoulos G, and Totlis T. Risk analysis of periprosthetic knee joint infection (PJI) in total knee arthroplasty after preoperative corticosteroid injection: a systematic review: A study performed by the Early-Osteoarthritis group of ESSKA-European Knee Associates section. *Arch Orthop Trauma Surg*. 2023;143(5):2683-2691.

Albanese J, Feltri P, Boffa A, Werner BC, Traina F, and Filardo G. Infection Risk Increases After Total Hip Arthroplasty Within 3 Months Following Intra-Articular Corticosteroid Injection. A Meta-Analysis on Knee and Hip Arthroplasty. *J Arthroplasty*. 2023;38(6):1184-1193.

Rhode DT, Siegel MA, Volchenko E, Patetta MJ, Schwarzman GR, Chmell SJ, and Gonzalez MH. Do Intra-articular Corticosteroid Injections Prior to Total Knee Arthroplasty Increase Postoperative Complication Rates: A Retrospective Review. *J Knee Surg*. 2023;36(3):231-235.